

Spectrum of Schrödinger field in a noncommutative magnetic monopole

D. Karabali^a, V.P. Nair^b and A.P. Polychronakos^{c,d1}

*a. Department of Physics and Astronomy
Lehman College of the CUNY, Bronx, NY 10468, USA*

*b. Physics Department, City College of the CUNY
New York, NY 10031, USA*

*c. Physics Department, The Rockefeller University
New York, NY 10021, USA*

*d. Physics Department, University of Ioannina
45110 Ioannina, Greece*

Abstract

The energy spectrum of a nonrelativistic particle on a noncommutative sphere in the presence of a magnetic monopole field is calculated. The system is treated in the field theory language, in which the one-particle sector of a charged Schrödinger field coupled to a noncommutative $U(1)$ gauge field is identified. It is shown that the Hamiltonian is essentially the angular momentum squared of the particle, but with a nontrivial scaling factor appearing, in agreement with the first-quantized canonical treatment of the problem. Monopole quantization is recovered and identified as the quantization of a commutative Seiberg-Witten mapped monopole field.

¹On leave from Theor. Physics Dept., Uppsala University, Sweden.
E-mail: karabali@alpha.lehman.cuny.edu, vpn@sci.ccny.cuny.edu, poly@teorfys.uu.se

1 Introduction

Noncommutative spaces arise in string theory and in the matrix model description of M-theory [1, 2, 3]. Gauge and field theories on such spaces have been extensively studied. For recent reviews, see [4, 5, 6].

The quantum mechanics of particles on noncommutative spaces has also been considered in some detail [7, 8, 9, 10, 11, 12, 13, 14]. In particular, the problem of a charged particle on the noncommutative plane [8] and the noncommutative sphere [7, 8] in an external constant magnetic field has been analyzed using the canonical approach, in which spatial noncommutativity relations complement the usual quantum mechanical commutation relations.

In field theory, particles emerge as field quanta. It is, therefore, of interest to recover the quantum mechanics of (non-interacting) particles as excitations of fields on noncommutative spaces with quadratic action. This would amount to obtaining and solving an equation on a noncommutative space for a Schrödinger-like field coupled to electromagnetic potentials. In [8] it was described how this can be done for the noncommutative plane, and was pointed out that the two approaches (canonical and field theory) are equivalent.

For the noncommutative sphere with a constant magnetic field, however, only the canonical approach was carried out. Although the canonical structure of the problem is quite clear, certain subtleties arise when one considers the limit in which the radius of the sphere goes to infinity, in relation to the correspondence with the planar results. Specifically, the Hamiltonian of the particle is not simply the square of the angular momentum, as one may naively expect; a magnetic field-dependent factor must be introduced in order to recover the correct planar limit. Further, monopole quantization (which is a purely group theoretic effect) does not imply the quantization of the strength of the magnetic field itself B , but rather of an effective magnetic field $\tilde{B} = B/(1 - \theta B)$. Both results clearly call for an explanation.

In this letter we give the field-theory treatment of the problem of the particle on the noncommutative sphere, by coupling a charged scalar field to a noncommutative monopole gauge field. The Hamiltonian is simply defined as the covariant kinetic term of this field. In the process, the scaling factor and the modified monopole quantization naturally appear and the correct planar limit is obtained, including even the zero-point energy of the ground state.

2 Review of $U(1)$ gauge theory on the noncommutative sphere

The noncommutative (NC) or ‘fuzzy’ sphere has been introduced a while ago [15] and field theories on this space have been studied rather extensively [16, 17, 18, 19]. We give here a self-contained exposition of the basic notions of gauge theory on the NC sphere using the operator (matrix) formulation.

We shall consider a noncommutative sphere defined in terms of the three coordinates x_i , satisfying

$$[x_i, x_j] = i \frac{\theta}{r} \epsilon_{ijk} x_k \quad (1)$$

θ and r are real parameters which, without loss of generality, can be chosen positive. (A negative θ/r corresponds to a parity transformation of the above space.) They will shortly be identified with the noncommutativity parameter and the radius of the sphere.

Clearly $R_i = (r/\theta)x_i$ satisfy the $SU(2)$ algebra. We shall choose the R_i to be in an irreducible representation of spin j . R_i and x_i become matrices of dimension $N = 2j + 1$. Requiring x_i^2 to be the square radius of space, namely r^2 , fixes the scaling of θ

$$\theta = \frac{r^2}{\sqrt{j(j+1)}} \quad (2)$$

Near the “north pole” (that is, for states near the highest spin state of the representation) we have $x_3 \simeq r$, so $[x_1, x_2] \simeq i\theta$. We can, therefore, identify θ above as the noncommutativity parameter of the space. Slightly different definitions with the same scaling as (2) for $j \rightarrow \infty$ are also acceptable. The definition

$$\theta' = \frac{r^2}{j + \frac{1}{2}} = \frac{4\pi r^2}{2\pi N} \quad (3)$$

in particular, identifies $2\pi\theta'$ as the elementary area on the sphere corresponding to each of the N states of the model, therefore making the allusion to phase space quantum mechanics clearer. We shall keep (2) as our definition of θ .

The two parameters r and j , or, equivalently, r and θ , define the model. Both the commutative sphere and the noncommutative plane are obtained as the limit $j \rightarrow \infty$ but with different scalings of r . Taking r fixed, $\theta \rightarrow 0$, that is, $j \rightarrow \infty$, recovers the commutative sphere, while taking $r \rightarrow \infty$, θ fixed, that is, $j \rightarrow r^2/\theta$ recovers the noncommutative plane.

We shall consider field theory on the noncommutative sphere with a commutative time. Scalar fields on the NC sphere correspond to arbitrary $N \times N$ matrices whose elements are functions of time. Such matrices can always be expanded in terms of monomials of x_i with any fixed ordering (such as the symmetric ordering). So they can be thought of as functions of the noncommutative coordinates of the sphere. The fact that the Casimir $x^2 = r^2$ is fixed reduces the number of independent functions in the same way as in the commutative sphere. The fact that x_i are of rank N , on the other hand, and therefore x_i^N can be reduced to lower-order polynomials of the x_i is a purely nonperturbative effect on the NC sphere and amounts to a cutoff to the possible ‘Fourier modes’ (that is, spherical harmonics) of the allowed functions, reflecting the finiteness of the Hilbert space of these functions.

The matrices $R_i = (r/\theta)x_i$ satisfy

$$[R_i, x_j] = i\epsilon_{ijk} x_k \quad (4)$$

Therefore, the adjoint action (commutator) of R_i on any function ψ of x_i generates spatial rotations of the x_i and can be identified with the angular momentum L_i :

$$L_i(\psi) \equiv [R_i, \psi] \quad (5)$$

Correspondingly, in analogy with the operator formulation of the NC plane, the operators

$$\partial_i = -\frac{i}{\theta} x_i = -\frac{i}{r} R_i \quad (6)$$

can be identified with partial derivatives. Integration over the NC sphere amounts to taking the trace with respect to the matrix space:

$$\int \psi = \frac{4\pi r^2}{N} \text{Tr} \psi \quad (7)$$

The prefactor in (7) is inserted to achieve compatibility with the proper commutative measure of integration; it ensures, for instance, that the area of the sphere is recovered as $\int 1 = 4\pi r^2$.

Gauge fields are introduced by deforming the above derivative operators into covariant derivatives

$$D_i = \partial_i + iA_i, \quad D_0 = \partial_0 + iA_0 \quad (8)$$

where A_i and A_0 are time-dependent hermitian $N \times N$ matrices. The A_i play the role of the three spatial components of the gauge field in the ambient three dimensional space of the sphere. In the above, ∂_0 is an ordinary differential operator while ∂_i are $N \times N$ matrices. The D_i are, thus, arbitrary time-dependent anti-hermitian matrices. Gauge transformations become conjugations of D_i and D_0 by the same arbitrary $U(N)$ matrix and correspond to a time-dependent change of basis for the N states on which the above matrices act.

The gauge field on a sphere has only 2 independent spatial components, since the radial component is missing. In the commutative case this is expressed by the relation

$$x_i A_i = 0 \quad (9)$$

which is also supplemented by the relation

$$x_i L_i = 0 \quad (10)$$

expressing the fact that L_i are derivatives along the surface of the sphere and do not act in the radial direction. To find the generalization of the above formulae in the NC case we note that from the covariant D_i we can construct new covariant coordinate fields $X_i = i\theta D_i$. In the absence of gauge fields, X_i are the undeformed coordinates x_i . In general, they can be interpreted as the NC coordinates of a membrane in the matrix regularization. The absence of a third component of the gauge field corresponds to the absence of radial excitations of the membrane; the membrane should remain spherical, that is,

$$X_i^2 = r^2 \quad (11)$$

In terms of the covariant derivative matrices D_i , this becomes

$$D_i^2 = -\frac{r^2}{\theta^2} = -\frac{j(j+1)}{r^2} \quad (12)$$

The above relation is obviously gauge invariant and replaces (9) in the NC case. Expressing $D_i = \partial_i + iA_i = -(i/\theta)x_i + iA_i$ we obtain

$$x_i A_i + A_i x_i - \theta A_i^2 = 0 \quad (13)$$

which becomes (9) in the limit $\theta \rightarrow 0$, where x_i and A_i commute. (10), on the other hand, takes the form

$$x_i L_i(\psi) + L_i(\psi) x_i = x_i [R_i, \psi] + [R_i, \psi] x_i = 0 \quad (14)$$

which, upon putting $R_i = (r/\theta)x_i$ and using $x_i^2 = r^2$, becomes an identity.

Since $[\partial_i, \partial_j] - \frac{1}{r}\epsilon_{ijk}\partial_k = 0$, the field strength F_{ij} is defined as

$$iF_{ij} = [D_i, D_j] - \frac{1}{r}\epsilon_{ijk}D_k \quad (15)$$

For $D_i = \partial_i$, the above definition gives $F_{ij} = 0$, as it should. F_{ij} is obviously gauge covariant under a gauge transformation,

$$D_i \rightarrow U D_i U^{-1} \implies F_{ij} \rightarrow U F_{ij} U^{-1} \quad (16)$$

In the planar limit $r \rightarrow \infty$, $iD_3 \rightarrow r/\theta$, iF_{ij} goes to $[D_i, D_j] + (i/\theta)$ which is the standard planar expression. The electric components of the field strength F_{0i} are, as usual,

$$F_{0i} = -i[D_0, D_i] = \partial_0 A_i - [D_i, A_0] \quad (17)$$

From F_{ij} we can define a magnetic field

$$iB_i = i\frac{1}{2}\epsilon_{ijk}F_{jk} = \epsilon_{ijk}D_j D_k - \frac{1}{r}D_i \quad (18)$$

A gauge covariant magnetic strength can now be defined in analogy to the commutative one as

$$B = \frac{X_i}{r}B_i = i\frac{\theta}{r}D_i B_i = \frac{\theta}{r}\epsilon_{ijk}D_i D_j D_k + \frac{1}{\theta}, \quad (19)$$

with a corresponding gauge invariant total flux over the sphere

$$\Phi = \int B = \frac{4\pi r\theta}{N}\epsilon_{ijk}\text{Tr}(D_i D_j D_k) + \frac{4\pi r^2}{\theta} \quad (20)$$

Note, however, that the above total flux is neither topologically conserved nor an integer multiple of 2π . The ‘correction’ $4\pi r^2/\theta = 4\pi\sqrt{j(j+1)}$ is almost $2\pi N$ and an appropriate definition of θ (different from either (2) or (3) above) would make it exactly $2\pi N$.

3 Monopole fields on the noncommutative sphere

The gauge field of a magnetic monopole at the center of the sphere would correspond to

$$B_i = \frac{B}{r} X_i = \frac{i\theta B}{r} D_i \quad (21)$$

with B a constant number. This is the gauge-invariant counterpart of the corresponding relation for a commutative abelian monopole field. Using (18) the above relation becomes

$$[D_i, D_j] = \frac{1 - \theta B}{r} \epsilon_{ijk} D_k \quad (22)$$

For $\theta B \neq 1$, this means that $rD_i/(1 - \theta B)$ satisfy the $SU(2)$ algebra. If the D_i are irreducible (that is, they cannot be brought to a block-diagonal form with an appropriate gauge transformation), since they are of dimension $N = 2j + 1$, they have to be proportional to ∂_i or x_i (or any gauge transformation of the above). Due to relation (12) which fixes their overall scale, we conclude that D_i have to be equal to ∂_i which implies $B = 0$. For $\theta B = 1$, on the other hand, the above relation implies that all D_i commute, and they are certainly completely reducible.

The only possibility, therefore, is to have reducible D_i . This means, however, that as far as the action of covariant derivatives is concerned, space has decomposed into non-communicating components, corresponding to the blocks. In the membrane picture, the membrane represented by the D_i consists of several concentric spherical components of the same radius without any connection between them. In the degenerate case of one-dimensional blocks (corresponding to $\theta B = 1$) the corresponding membranes are single points at the surface of a sphere of radius r . As a result, the dynamics of a scalar wavefunction corresponding to reducible D_i 's decomposes into non-communicating sectors, each corresponding to a different sphere. For an irreducible sector of dimension $M = 2k + 1$, the commutation relation (22) and normalization condition (12) imply

$$(1 - \theta B)^2 = \frac{j(j+1)}{k(k+1)} \equiv \gamma^2 \quad (23)$$

where we defined the dimensionless factor $\gamma = 1 - \theta B$. Therefore, the monopole field B in each of these sectors is determined by the dimensionality of the sector. There is no need to choose the same monopole field B in each of these sectors since they are not communicating, and we can restrict ourselves in one of these sectors of dimension $M = 2k + 1$ corresponding to a desired B field. This is the central observation, which will be further justified when we analyze the Schrödinger equation in the next section.

In conclusion, for a monopole field we can write the D_i in a block-diagonal form: the upper component D_i^u , of dimensionality $M = 2k + 1$, is of the form

$$D_i^u = \frac{1 - \theta B}{r} K_i = \frac{\gamma}{r} K_i \quad (24)$$

with K_i irreducible $SU(2)$ matrices of spin k . In the above we used relation (23) for the monopole field B . D_i^u obviously satisfies (22) and (12). The lower component D_i^ℓ , of dimensionality $M' = N - M$, corresponds to the decoupled part and can be any set of matrices satisfying (22) and (12). It could be, for instance, a diagonal part with entries $-(i/\theta)z_{i,m}$, $m = 1, \dots, M'$. This gauge field is of the projector type [20, 21] and its field strength corresponds to M' ‘fuzzy Dirac strings’ at positions z_i on the sphere which can be thought of as bringing in the flux [22]. Alternatively, the lower component could also be an irreducible space of the form (24) for some $k' = j - k - \frac{1}{2}$, corresponding to a disconnected fuzzy sphere with the ‘complementary’ magnetic field.

It is clear from the above that there is a constraint on the dimensionalities M and M' (summing to N) and a ‘monopole quantization’ corresponding to the fact that $2k$ is always an integer. This, however, does not translate into the quantization of the conventional flux Φ as defined in (20). To get a feeling of what this monopole quantization corresponds to, we solve (23) for B and calculate the quantity $\tilde{B} = B/(1 - \theta B)$:

$$\tilde{B} = \frac{B}{1 - \theta B} = \frac{1}{r^2} \left(\pm \sqrt{k(k+1)} - \sqrt{j(j+1)} \right) \quad (25)$$

In either the commutative sphere limit or the noncommutative plane limit, $j \rightarrow \infty$. If we want the total flux $\tilde{\Phi}$ of the field $\tilde{B}$ through the surface of the sphere to remain finite, we must choose the positive sign in (25) and take the difference $k - j$ to remain finite. In that limit we obtain

$$\tilde{\Phi} = 4\pi r^2 \tilde{B} = 2\pi \cdot 2(k - j) \quad (26)$$

Thus the flux $\tilde{\Phi}$ is quantized in integer multiples of 2π , which is the standard monopole quantization.

That $\tilde{B}$ rather than B enters the quantization condition is not so surprising. It is $\tilde{B}$, for instance, which determines the density of states for a charged particle in each Landau level on the noncommutative plane [8]. A related analysis on the noncommutative two-torus also revealed the special role of $\tilde{B}$ [10]. Finally, the Seiberg-Witten transformation [3], which maps a noncommutative gauge field to a corresponding commutative one, actually maps a constant noncommutative field B to a commutative one $\tilde{B} = B/(1 - \theta B)$. Since the gauge transformation properties of the field $\tilde{B}$ are the usual commutative abelian ones, we expect monopole quantization to hold for $\tilde{B}$.

One last remark is in order. Just as we have taken $k < j$ in the above, we could formally take $k > j$ to obtain the opposite monopole number. This cannot, of course, be obtained through a reducible representation of the D_i ’s in the original space, but it can be obtained as a representation in a larger space. The point remains, in each case, that a monopole charge in a NC sphere corresponds to *two* NC spheres, one with spin j representing the uncharged sphere and one with different spin k representing the charged sphere, with their difference fixed by the monopole number.

We conclude by mentioning that monopoles in the NC sphere and the associated Dirac operators have been studied in [23, 24] using sections of bundles or partial isometries approach. Our analysis in this section serves as a more conventional treatment in the standard NC field theory language.

4 Charged particle field on the NC sphere with a monopole

We can now address the issue of the quantum states and energies of a charged particle on the NC sphere with a magnetic monopole field. We shall consider time-independent magnetic field with a monopole charge $n = 2(k - j)$ constructed as above and take $A_0 = 0$.

In noncommutative gauge theory, matter fields can only be in the fundamental (F), antifundamental ($\overline{F}$) or adjoint (A) representation. The origin of this restriction is very simple and has to do with the possible couplings of a field to the gauge covariant operators D_μ . A matter field ψ is represented as a matrix in noncommutative space. The only possible couplings to the operators D_μ which preserve the matrix structure are matrix multiplications. Considering, then, kinetic terms that reduce to the proper derivative (commutator) form $\partial_i(\phi) = [\partial_i, \phi]$ in the zero gauge field limit, we have the possibilities,

$$F: D_i(\psi) = D_i\psi - \psi\partial_i, \quad \psi \rightarrow U\psi \quad (27)$$

$$\overline{F}: D_i(\psi) = \partial_i\psi - \psi D_i, \quad \psi \rightarrow \psi U^{-1} \quad (28)$$

$$A: D_i(\psi) = D_i\psi - \psi D_i, \quad \psi \rightarrow U\psi U^{-1} \quad (29)$$

In the $U(1)$ case, the adjoint matter field would decouple in the commutative limit and does not correspond to a charged field. The fundamental and antifundamental cases correspond to fields with charges $+1$ and -1 in natural units. Their treatment and corresponding spectra are essentially identical. For our analysis we shall choose the antifundamental case, to conform with the conventions of [8] and to make the monopole field above (with monopole number $n = 2(k - j) < 0$) a positively charged one.

We consider a Lagrangian for ψ of the Schrödinger type:

$$\mathcal{L} = \frac{4\pi r^2}{N} \text{Tr} \left(i\psi^\dagger \dot{\psi} + \frac{1}{2} D_i(\psi)^\dagger D_i(\psi) \right) \quad (30)$$

As usual, the one-particle sector of the field theory defined by this Lagrangian is described by a wavefunction ψ which satisfies the Schrödinger equation implied by (30). The corresponding energy eigenvalue problem is

$$H(\psi) = -\frac{1}{2} D_i^2(\psi) = E\psi \quad (31)$$

where $D_i(\psi)$ is defined in (28).

In the covariant derivative (28) above, ∂_i are irreducible $N \times N$ matrices, but D_i are reducible block-diagonal matrices. As a result, $D_i(\psi)$ decouples into two non-mixing components. We decompose ψ into two rectangular components

$$\psi = (\phi | \phi') \quad (32)$$

where ϕ is a $M \times N$ matrix, representing the first M columns of ψ , while ϕ' is a $M' \times N$ matrix representing the remaining M' columns. The covariant derivative of ψ then takes the form

$$iD_i(\psi) = \left(iD_i^u(\phi) \mid iD_i^\ell(\phi') \right) = \frac{1}{r} (R_i\phi - \gamma\phi K_i \mid R_i\phi' - \gamma'\phi' K_i') \quad (33)$$

So $D_i(\psi)$ breaks into two decoupling components, the one of interest for our problem ϕ and the complementary one ϕ' . The kinetic energy term of ψ , $-\frac{1}{2}D_i^2(\psi)$, appearing in the Schrödinger equation, similarly decomposes into two decoupling components, one for ϕ and one for ϕ' . The particle could be either in the ‘upper’ sphere, with wavefunction ϕ or in the ‘lower’ one with wavefunction ϕ' . Although we could consider linear superpositions of these states, there is nothing that could mix the two components and therefore we may choose as a superselection rule $\phi' = 0$. This justifies the restriction to the upper component of space alone, as argued in the previous section. On this component we have

$$iD_i(\phi) = \frac{1}{r}R_i\phi - \frac{\gamma}{r}\phi K_i \quad (34)$$

We note that the matrix R_i acts on ϕ on the left, while the matrix $-K_i$ acts on the right. As a result, when acting on ϕ , R_i and K_j commute. Note, further, that $-K_i$ when acting on the right of ϕ , satisfy $SU(2)$ commutation relations. We can therefore think of the $M \times N$ degrees of freedom in the matrix ϕ as spanning the tensor product of two spaces V_R and V_K on which R_i and K_i act, V_R being the N states of the spin- j representation of $SU(2)$ carried by R_i and V_K being the M states of the spin- k representation carried by K_i . So, the matrix elements ϕ_{mn} (which parametrize the quantum state of the particle) go over to a state $|\phi\rangle$

$$|\phi\rangle = \sum_{n,m} \phi_{mn} |k, m\rangle \otimes |j, n\rangle \quad (35)$$

with $|j, n\rangle$, $n = -j, \dots, j$, being the states in a spin- j multiplet. Altogether, the covariant derivative $D_i(\phi)$ can be written as

$$iD_i|\phi\rangle = \frac{1}{r}(R_i + \gamma K_i)|\phi\rangle \quad (36)$$

where, now, K_i is the operator acting on $|\phi\rangle$ corresponding to $-K_i$ acting by right-multiplication on the matrix ϕ .

We have therefore made contact with the setting of [8]. The R_i and K_i above are the corresponding operators defined there in the canonical framework. The new element, however, is the appearance of the scaling factor γ in the covariant derivative above. We also note that the covariant angular momentum operators on $|\phi\rangle$, defined as the operators J_i which satisfy $SU(2)$ commutation relations and commute as in (4) with x_i , are

$$J_i(\phi) = R_i\phi - \phi K_i, \quad J_i|\phi\rangle = (R_i + K_i)|\phi\rangle \quad (37)$$

as in [8]. This can also be identified from the invariance of the Schrödinger action. Consider the transformations generated by

$$\delta\phi = i\omega_j(R_j\phi - \phi K_j) \quad (38)$$

with ω_j time-independent real numbers. The covariant derivative (34) on ϕ transforms as

$$\delta D_i(\phi) = i\omega_j \{R_j D_i(\phi) - D_i(\phi) K_j - \epsilon_{ijk} D_k(\phi)\} \quad (39)$$

It can be checked that the Schrödinger Lagrangian (30) is invariant under the above transformation. The Noether charge derived from this symmetry is

$$Q = \omega_i Q_i = \omega_i \frac{4\pi r^2}{N} \text{Tr} [\phi^\dagger (R_i\phi - \phi K_i)] \quad (40)$$

and corresponds to the expectation value of the operator J_i as defined in (37).

We can finally calculate the spectrum of the Hamiltonian $H = -\frac{1}{2}D_i^2$. On states $|\phi\rangle$ the above Hamiltonian takes the form

$$H = \frac{1}{2r^2}(R_i + \gamma K_i)^2 = \frac{1}{2r^2} \left(\gamma(R_i + K_i)^2 + (1 - \gamma)R_i^2 - \gamma(1 - \gamma)K_i^2 \right) \quad (41)$$

Using the expressions for the Casimirs $R_i^2 = j(j+1) = r^4/\theta^2$ and $K_i^2 = k(k+1) = \gamma^{-2}R_i^2$ we finally obtain

$$H = \frac{\gamma}{2r^2} \left\{ J^2 - \left(\frac{Br^2}{\gamma} \right)^2 \right\} \quad (42)$$

where we used (23) to express

$$\frac{Br^2}{\gamma} = \frac{r^2}{\theta} \left(\frac{1}{\gamma} - 1 \right) = \pm \sqrt{k(k+1)} - \sqrt{j(j+1)} \quad (43)$$

This is exactly the Hamiltonian identified in [8] (up to a zero-point shift), with the nontrivial scaling factor γ naturally appearing.

What is more, the above Hamiltonian even has the correct zero-point energy as compared with the planar limit. To see this, note that $J_i = R_i + K_i$ is simply the sum of two commuting $SU(2)$ operators and its Casimir J^2 takes values $\ell(\ell+1)$ with $\ell = |j-k|, \dots, j+k$. The total angular momentum has a lowest value equal

to half the monopole number, as expected, and a highest value $j + k$ reflecting the finiteness of the NC Hilbert space. Its ground state is $J_0^2 = |j - k|(|j - k| + 1)$. In the limit of the NC plane, both j and k go to infinity; by (2), j becomes $r^2/\theta - \frac{1}{2}$ in this limit, while, by (23), k becomes $r^2/(\gamma\theta) - \frac{1}{2}$. Their difference $j - k$ then becomes (taking γ positive)

$$|j - k| \rightarrow \frac{r^2}{\theta} \left| 1 - \frac{1}{\gamma} \right| = \frac{|B|r^2}{\gamma} \quad (44)$$

Putting everything together, the ground state of H in (42) becomes

$$E_0 = \frac{|B|}{2} \quad (45)$$

which is the correct zero-point energy of the lowest Landau level. The rest of the spectrum also matches, as demonstrated in [8].

5 Adjoint matter field

One can similarly analyze the spectrum for an adjoint matter field. The energy eigenvalue problem is the same as in (31), but with $D_i(\psi)$ defined as in (29). ψ is decomposed into

$$\psi = \begin{pmatrix} \phi_1 & \phi_2 \\ \phi_3 & \phi_4 \end{pmatrix} \quad (46)$$

where ϕ_1 is a $M \times M$ matrix, ϕ_2 is a $M \times M'$ matrix, ϕ_3 is a $M' \times M$ matrix and ϕ_4 is a $M' \times M'$ matrix. The covariant derivative is of the form

$$D_i \psi = \begin{pmatrix} [D_i^u, \phi_1] & D_i^u \phi_2 - \phi_2 D_i^l \\ D_i^l \phi_3 - \phi_3 D_i^u & [D_i^l, \phi_4] \end{pmatrix} \quad (47)$$

The corresponding kinetic energy term decomposes into four decoupled components and we can write

$$H = -\frac{1}{2} (D_L + D_R)^2 \quad (48)$$

where for the four ϕ 's in ψ we have the assignments

$$\begin{aligned} \phi_1 & : \quad D_L = \frac{i\gamma K}{r}, \quad D_R = \frac{i\gamma K}{r} \\ \phi_2 & : \quad D_L = \frac{i\gamma K}{r}, \quad D_R = \frac{i\gamma' K'}{r} \\ \phi_3 & : \quad D_L = \frac{i\gamma' K'}{r}, \quad D_R = \frac{i\gamma K}{r} \\ \phi_4 & : \quad D_L = \frac{i\gamma' K'}{r}, \quad D_R = \frac{i\gamma' K'}{r} \end{aligned} \quad (49)$$

ϕ_1 can be expanded as $\phi_1 = \sum \phi_{1nm} |k, n\rangle \otimes |k, m\rangle$ with the energy eigenvalues

$$E_1 = \frac{\gamma^2}{2r^2} \ell(\ell + 1) \quad (50)$$

$l = 0, 1, \dots, 2k$. As $j, k \rightarrow \infty$, this coincides with the spectrum of a free uncharged particle on the sphere. In the commutative limit the adjoint field has no charge and so does not couple to the monopole. The above result is as expected. Similar results hold for ϕ_4 as $j, k' \rightarrow \infty$. For $k' = j - k - 1/2$ finite and $j, k \rightarrow \infty$, the ϕ_4 field has the spectrum

$$E_4 = \frac{\gamma'^2}{2r^2} \ell(\ell + 1) \quad (51)$$

$l = 0, 1, \dots, 2k'$. $E_4 \sim j(j + 1)$ at large j and the number of states remains finite as $j, k \rightarrow \infty$, with k' fixed. (Presumably these modes become nonpropagating as $j \rightarrow \infty$.) The “monopole field” seen by the upper components is proportional to $j - k$, the field seen by the lower components is proportional to $j - k'$. Therefore if either is fixed and we take $j \rightarrow \infty$, the field seen by the other becomes very large and this leads to the above behavior of energy eigenvalues.

For the ϕ_2 field we have

$$\begin{aligned} E_2 &= \frac{(\gamma K + \gamma' K')^2}{2r^2} \\ &= \frac{j(j + 1)}{2r^2} \left[2 - \sqrt{\frac{k(k + 1)}{k'(k' + 1)}} - \sqrt{\frac{k'(k' + 1)}{k(k + 1)}} + \frac{(K + K')^2}{\sqrt{k(k + 1)k'(k' + 1)}} \right] \end{aligned} \quad (52)$$

For finite monopole number, the lowest value of E_2 diverges as $j \rightarrow \infty$, but the limit $j, k, k' \rightarrow \infty$ with $(k - k')$ fixed can give finite energies. A similar statement holds for the ϕ_3 field as well.

The adjoint fields behave in some respects like combinations of the antifundamental representations suitably on the upper and lower spheres. To see this, consider a field ψ which transforms as $\phi \otimes \phi'$, where ϕ, ϕ' are in the antifundamental representations as in section 4, ϕ on the upper sphere and ϕ' on the lower sphere. ψ may be expanded as

$$\psi \sim \phi \otimes \phi' = \sum \psi_{mnm'n'} |k, m\rangle \otimes |j, n\rangle \otimes |k', m'\rangle \otimes |j, n'\rangle \quad (53)$$

The action of the covariant derivative is, by using formula (33),

$$iD_i(\psi) = (R_i + \gamma K_i + \gamma' K'_i) \psi \quad (54)$$

R_i acts on $|j, n\rangle$ and $|j, n'\rangle$, both being spin- j representations. The singlet combination in $\phi \otimes \phi'$ with $R_i = 0$ behaves like ϕ_2 . One can think of the other components of the adjoint in a similar way.

6 Two subtleties

In the analysis of the previous sections we glossed over two subtle points, which we wish to clarify here. They have to do with the $\pm$ sign in (25,43) and with the critical magnetic field $\theta B = 1$.

Equation (23) for the magnetic field has two solutions, corresponding to $\gamma > 0$ and $\gamma < 0$. This is the origin of the double signs appearing in (25,43). Thus, the same values of j, k represent two values of the magnetic field, one subcritical $B_- < 1/\theta$ and one overcritical $B_+ > 1/\theta$, corresponding to $\gamma = \pm|\gamma|$ and related by

$$B_- + B_+ = \frac{2}{\theta} \quad (55)$$

The matrices R_i, K_i describing the model are the same and the only difference is the sign of the factor γ appearing in the expressions for $D_i(\phi)$ and H . To see how the two cases are imbedded in the Hilbert space, notice that the Hamiltonian has a prefactor γ . Thus, if $\gamma > 0$ (corresponding to $B_- < 1/\theta$) the ground state of H corresponds to the lowest eigenvalue of J^2 (as assumed in the last paragraph of section 5). If, however, $\gamma < 0$ (corresponding to $B_+ > 1/\theta$) the ground state corresponds to the highest eigenvalue of J^2 . The analysis of the previous section still goes through, with $j + k$, now, defining the ground state, and the energy levels in terms of B are as before. We conclude that the energy spectrum for the two values of the magnetic field $B_{\pm}$ is essentially the same but reversed, with the states at each end of the spectrum mapping to two different magnetic fields in the planar limit, as explained in [8]. Note, also, that for subcritical fields the angular momentum for excited states increases, as expected, while for overcritical fields the angular momentum *decreases* with energy. (A similar effect for the planar case was pointed out in [12].)

There is a potential confusion with the critical field $B = 1/\theta$ corresponding to $\gamma = 0$. From (23) it appears that $\gamma = 0$ corresponds to the limit $k \rightarrow \infty$. On the other hand, from equation (22) it appears that $\gamma = 0$ corresponds to the completely reducible case in which we should choose an irreducible $D_i = -(i/\theta)z_i$ with z_i a real vector squaring to r^2 . We now clarify the relationship between these two cases.

There is a related issue in the planar case. In the first-quantized picture, the algebra of observables for $\theta B = 1$ becomes reducible. An irreducible Hilbert space would put $x_i + \theta\epsilon_{ij}p_j$ ($i, j = 1, 2$) equal to some numbers z_i , since they become Casimirs at the critical value of B , and would correspond to a single Heisenberg copy of the algebra of the x_i 's alone. The limit of the Hilbert space as $B \rightarrow 1/\theta$, on the other hand, contains an infinite number of copies of the irreducible space, embedded into the direct product of two Heisenberg representations. An irreducible component in this space corresponds to picking one state from each Landau level, centered around the spatial point z_i corresponding to the value of the Casimir.

On the sphere, the algebra of observables is always irreducible and, a priori, there appears to be no mechanism for recovering the irreducible planar Hilbert space for $B = 1/\theta$. In fact, the reducibility of (22) achieves just that. The representation

obtained as the limit $k \rightarrow \infty$ maps to the limit $B \rightarrow 1/\theta$ in the planar case, while the representation $D_i = -(i/\theta)z_i$ maps to the irreducible planar Hilbert space. To see this, note that the wavefunction ϕ for this choice of D_i becomes a single copy of the spin- j multiplet, on which D_i acts as

$$iD_i(\phi) = \left(\frac{1}{r}R_i\phi - \frac{z_i}{\theta} \right) \phi \quad (56)$$

The Hamiltonian becomes

$$H = \frac{1}{2} \left(\frac{1}{r}R_i - \frac{z_i}{\theta} \right)^2 = \frac{r^2}{\theta^2} - \frac{z_i R_i}{\theta r} \quad (57)$$

where we used $R_i^2 = r^4/\theta^2$ and $z_i^2 = r^2$. The matrix $\hat{R} = (z_i/r)R_i$ is clearly a spin- j $SU(2)$ generator and has eigenvalues $\hat{R} = j - n$, $n = 0, 1, \dots, 2j$. So the energy levels are

$$E_n = \frac{n}{\theta} + \frac{r^2}{\theta^2} - \frac{j}{\theta} \quad (58)$$

each with degeneracy one.

In the limit $r \rightarrow \infty$, j becomes $r^2/\theta - \frac{1}{2}$. So the energy levels become

$$E_n = \frac{1}{\theta} \left(n + \frac{1}{2} \right) \quad (59)$$

These are exactly the Landau energy levels for $B = 1/\theta$, with one state per level, as in the irreducible planar Hilbert space. For these states the expectation value of R_i is $\langle R_i \rangle = (z_i/r)\langle \hat{R} \rangle$ and we get

$$\langle x_i \rangle = \frac{\theta}{r} \langle R_i \rangle = \frac{\theta z_i}{r^2} (j - n) \rightarrow z_i - \frac{\theta z_i}{r^2} \left(n + \frac{1}{2} \right) \quad (60)$$

So in the limit $r \rightarrow \infty$ we get $\langle x_i \rangle = z_i$, as in the planar case.

The reducibility of the system at $\theta B = 1$ can be lifted if we include the generators of spatial symmetries in the algebra of observables. In the planar case, the Casimirs $x_i + \theta \epsilon_{ij} p_j$ are not invariant under translations and rotations. That is, the translation and rotation generators cannot be expressed any more in terms of x_i and p_i , and independent generators for these translations have to be appended, rendering the algebra irreducible and calling for the full Hilbert space of two copies of the Heisenberg representation, as obtained in the $B \rightarrow 1/\theta$ limit. This is the situation closest resembling the sphere. The full algebra of x_i and J_i is irreducible. The reduced representation corresponding to (56) breaks rotational invariance (which would rotate the Casimirs z_i). (Note that (39) does not hold if $\delta\phi$ is as in (38) but $D_i\phi$ is as in (56).) Introducing appropriate J_i which also rotate the z_i calls for inclusion of representations with all possible z_i , reproducing the infinite-dimensional Hilbert space as obtained in the $k \rightarrow \infty$ limit.

In summary, the situation for $B = 1/\theta$ is similar between the plane and the sphere except for the fact that on the sphere we already have the J_i 's which can

make changes (rotations) in z_i . On the plane we have to enlarge the algebra of observables to incorporate shifts of z_i .

Let us also comment on the limit $k \rightarrow 0$, which corresponds to $B \rightarrow \pm\infty$. In that limit, the field ϕ becomes again a single copy of the spin- j multiplet, as in the reduced case $B = 1/\theta$ above. This could be surprising, since the two cases represent vastly different values of B . What happens in this case, in fact, is the well-known reduction of the Hilbert space in a strong magnetic field to the lowest Landau level. The dimensionality of the Hilbert space is $(2j+1)(2k+1)$. As $|B|$ becomes large, k decreases, so the dimensionality of the space decreases. The density of states per Landau level, on the other hand, increases linearly with $|\tilde{B}|$ and becomes θ^{-1} for large $|B|$. So the Hilbert space can accomodate fewer and fewer Landau levels and, at the limit $k \rightarrow 0$, only one level remains. The energy spectrum is quite different in the two cases. For $k \rightarrow 0$ all $2j+1$ states become degenerate with a divergent energy, while for the reduced $\theta B = 1$ case they reproduce an equidistant spectrum with spacing θ^{-1} .

In summary, the NC sphere-cum-monopole presents quite a fascinating ‘package’: it captures all the features and special limits of planar noncommutative theory, for two values of the magnetic field at a time, all encoded in a finite Hilbert space.

7 Conclusions

We have analyzed the problem of a charged particle on the NC sphere from the NC gauge field point of view. We identified the monopole sectors and obtained the spectrum of a charged Schrödinger field both for the fundamental and adjoint couplings. Our results are in complete agreement with the canonical analysis of the problem, and reproduce the nontrivial scaling factor of the Hamiltonian, as well as the zero-point shift in the energy required to obtain the correct correspondence with the planar limit. We also obtained the magnetic monopole quantization and identified it as the quantization of an equivalent commutative monopole charge obtained through the Seiberg-Witten transformation.

Some other issues of the dynamics of a charged particle on the NC sphere in the presence of a monopole, such as inclusion of spin, are under investigation.

Acknowledgements

We would like to thank B. Morariu for discussions. This work was supported in part by the National Science Foundation grants PHY-9970724 and PHY-0070883 and PSC-CUNY-32 awards.

References

- [1] T. Banks, W. Fischler, S. H. Shenker and L. Susskind, Phys. Rev. D **55**, 5112 (1997) [arXiv:hep-th/9610043].
- [2] A. Connes, M. R. Douglas and A. Schwarz, JHEP **9802**, 003 (1998) [arXiv:hep-th/9711162].
- [3] N. Seiberg and E. Witten, JHEP **9909**, 032 (1999) [arXiv:hep-th/9908142].
- [4] J. A. Harvey, arXiv:hep-th/0102076.
- [5] M. R. Douglas and N. A. Nekrasov, arXiv:hep-th/0106048.
- [6] R. J. Szabo, arXiv:hep-th/0109162.
- [7] V. P. Nair, Phys. Lett. B **505**, 249 (2001) [arXiv:hep-th/0008027].
- [8] V. P. Nair and A. P. Polychronakos, Phys. Lett. B **505**, 267 (2001) [arXiv:hep-th/0011172].
- [9] J. Gamboa, M. Loewe and J. C. Rojas, Phys. Rev. D **64**, 067901 (2001) [arXiv:hep-th/0010220]; J. Gamboa, M. Loewe, F. Mendez and J. C. Rojas, arXiv:hep-th/0106125.
- [10] B. Morariu and A. P. Polychronakos, Nucl. Phys. B **610**, 531 (2001) [arXiv:hep-th/0102157].
- [11] C. Duval and P. A. Horvathy, J. Phys. A **34**, 10097 (2001) [arXiv:hep-th/0106089].
- [12] S. Bellucci, A. Nersessian and C. Sochichiu, arXiv:hep-th/0106138.
- [13] P. M. Ho and H. C. Kao, arXiv:hep-th/0110191.
- [14] R. Iengo and R. Ramachandran, arXiv:hep-th/0111200.
- [15] J. Madore, Phys. Lett. **B263**, 245 (1991); Class. Quant. Grav. **9**, 69 (1992);
- [16] U. Carow-Watamura and S. Watamura, Commun. Math. Phys. **183**, 365 (1997) [arXiv:hep-th/9605003]; Commun. Math. Phys. **212**, 395 (2000) [arXiv:hep-th/9801195]; Int. J. Mod. Phys. A **13**, 3235 (1998) [arXiv:q-alg/9710034].
- [17] C. Klimcik, Commun. Math. Phys. **199**, 257 (1998) [arXiv:hep-th/9710153].
- [18] J. Gratus, arXiv:q-alg/9710014.
- [19] S. Iso, Y. Kimura, K. Tanaka and K. Wakatsuki, Nucl. Phys. B **604**, 121 (2001) [arXiv:hep-th/0101102].
- [20] A. P. Polychronakos, Phys. Lett. B **495**, 407 (2000) [arXiv:hep-th/0007043].

- [21] D. J. Gross and N. A. Nekrasov, JHEP **0010**, 021 (2000) [arXiv:hep-th/0007204]; JHEP **0103**, 044 (2001) [arXiv:hep-th/0010090].
- [22] K. Hashimoto and K. Krasnov, Phys. Rev. D **64**, 046007 (2001) [arXiv:hep-th/0101145].
- [23] H. Grosse and P. Presnajder, Lett. Math. Phys. **33**, 171 (1995); H. Grosse, C. Klimcik and P. Presnajder, Commun. Math. Phys. **178**, 507 (1996) [arXiv:hep-th/9510083].
- [24] S. Baez, A. P. Balachandran, S. Vaidya and B. Ydri, Commun. Math. Phys. **208**, 787 (2000) [arXiv:hep-th/9811169]; A. P. Balachandran and S. Vaidya, Int. J. Mod. Phys. A **16**, 17 (2001) [arXiv:hep-th/9910129].